CLAIMS



5

1. A method of labeling oligonucleotides, comprising:

- a) providing: i) a solid support-bound oligonucleotide comprising an amino group, ii) a bifunctional linker arm and iii) an activated label;
- b) reacting said solid support-bound oligonucleotide with said bifunctional linker arm to produce a support-bound, linker-oligonucleotide;
- c) reacting said support bound linker-oligonucleotide with said activated label to produce a labeled support-bound protected oligonucleotide.

2. The method of Claim 1 wherein said bifunctional linker arm is selected from a group consisting of the compounds listed in Tables 1, 2 and 3.

3. The method of Claim 1 wherein said activated label is selected from a group consisting of the compounds listed in Table 1, 2 and 3.

4. A method of labeling oligonucleotides, comprising:

- a) providing: i) a solid support-bound oligonucleotide comprising an amino group, ii) a bifunctional linker arm and iii) an activated label;
- b) reacting said solid support-bound oligonucleotide with said bifunctional linker arm to produce a support-bound, protected linker-oligonucleotide;
- c) deprotecting the amino group of said support-bound, protected linkeroligonucleotide to produce a support-bound deprotected linker-oligonucleotide, and;
- d) reacting said support-bound deprotected linker-oligonucleotide with said activated label to produce a labeled support-bound protected oligonucleotide.

10

15 12 12

j

20